

EVEN MORE CALCULUS

Text

Contents

Section

- 1 Derivatives of Trig Functions
- 2 Rates of Change
- 3 Integration by Substitution
- 4 Integration by Parts
- 5 First Order Differential Equations

Even More Calculus

Introduction

You should be familiar with the differentiation and integration of functions such as x^n , e^x and $\ln x$. In this unit the range of functions is expanded to include the differentiation and integration of trigonometric functions and two important methods of integration, namely substitution and by parts, are dealt with. Finally, there is an introduction to the important topic of differential equations.

1 Derivatives of Trig Functions

Intuitive ideas

The derivative of a general function $f(x)$ is given by

$$f'(x) = \lim_{h \rightarrow 0} \left\{ \frac{f(x+h) - f(x)}{h} \right\}$$


You are going to obtain the derivatives of $\sin x$, $\cos x$, and $\tan x$.

Remember that the derivative of a function at a given point is given by the **gradient** of that function at the given point.


Suppose you look at a sketch of the graph $\sin x$ and sketch in the tangents for values of x of

$$-\frac{3\pi}{2}, -\pi, -\frac{\pi}{2}, 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}$$


This is shown opposite.



These tangent values when plotted against x will look like the sketch opposite.



This suggests that the graph of the function for the tangent may be something like the sketch opposite so this sketch suggests that the derivative of $\sin x$ might be $\cos x$, but this illustration is certainly not a proof.



Using a calculator or computer The derivative of $\sin x$, working in radians, is given by

$$\frac{d}{dx}(\sin x) = \lim_{h \rightarrow 0} \left\{ \frac{\sin(x+h) - \sin x}{h} \right\}$$

Using a calculator or computer you can try some specific examples - using a set value for x and making h small.

For instance, setting $x = 0.2$ and $h = 0.00004$,

$$\begin{aligned} \frac{\sin(0.20004) - \sin(0.2)}{0.00004} &= \frac{0.198708533 - 0.198669331}{0.00004} \\ &= \frac{0.000039202}{0.00004} \\ &= 0.98005 \\ &= \cos(0.200083427) \end{aligned}$$

So, approximately

$$\frac{\sin(0.20004) - \sin(0.2)}{0.00004} \approx \cos(0.2)$$

We could try another small value of h , but this would still not be a proof.

Geometric proof

It is also possible to use the unit circle to demonstrate that, for example,

$$\frac{d}{dx}(\sin x) = \cos x$$


In the diagram, shown opposite, h is small, and

$$A'B' = \sin(x+h)$$

$$AB = \sin x$$

$$A'A = h \quad (\text{radius} \times \text{angle})$$

So $\sin(x+h) - \sin x \approx A'A$




Now assume that when h is very small the arc $A'A$ is approximately straight.

So for the 'triangle' $AA'P$,

$$\frac{\sin(x+h) - \sin x}{h} \approx \frac{A'P}{A'A} = \cos x$$

So, approximately, if h is very small

$$\frac{\sin(x+h) - \sin x}{h} \approx \cos x$$



Now, taking the limit as $h \rightarrow 0$, gives

$$\frac{d}{dx}(\sin x) = \cos x$$

In a similar way, we can show that

$$\frac{d}{dx}(\cos x) = -\sin x$$

To find the derivative of $\tan x$, formally you can use the quotient rule,

$$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{u'v - v'u}{v^2} \quad \left(u' = \frac{du}{dx}, v' = \frac{dv}{dx}\right),$$

giving

$$\begin{aligned} \frac{d}{dx}(\tan x) &= \frac{d}{dx}\left(\frac{\sin x}{\cos x}\right) \\ &= \frac{\frac{d}{dx}(\sin x) \times \cos x - \sin x \times \frac{d}{dx}(\cos x)}{(\cos x)^2} \\ &= \frac{\cos x \times \cos x - \sin x \times (-\sin x)}{(\cos x)^2} \\ &= \frac{(\cos x)^2 + (\sin x)^2}{(\cos x)^2} \\ &= \frac{1}{(\cos x)^2} \end{aligned}$$

This gives

$$\frac{d}{dx}(\tan x) = (\sec x)^2$$



Worked Example 1

Show that

$$\frac{d}{dx}\left(\frac{1}{\cot x}\right) = -\cos^2 x$$



Solution

$$\begin{aligned} \frac{d}{dx}(\cot x) &= \frac{d}{dx}\left(\frac{1}{\tan x}\right) \\ &= \frac{d}{dx}(\tan x)^{-1} \\ &= \frac{-1}{(\tan x)^2} \frac{d}{dx}(\tan x) \end{aligned}$$

$$= \frac{-1}{(\tan x)^2} \cdot (\sec x)^2$$

$$= \frac{(\cos x)^2}{(\sin x)^2} \cdot \frac{1}{(\cos x)^2} = \frac{-1}{(\sin x)^2} = \operatorname{cosec}^2 x$$

So the three crucial results are

$$\frac{d}{dx}(\sin x) = \cos x \quad \text{or} \quad y = \sin x \Rightarrow \frac{dy}{dx} = \cos x$$

$$\frac{d}{dx}(\cos x) = -\sin x \quad \text{or} \quad y = \cos x \Rightarrow \frac{dy}{dx} = -\sin x$$

$$\frac{d}{dx}(\tan x) = \sec^2 x \quad \text{or} \quad y = \tan x \Rightarrow \frac{dy}{dx} = \sec^2 x$$

and, inverting these results

$$\int \cos x \, dx = \sin x + c$$

$$\int \sin x \, dx = -\cos x + c$$

$$\int \sec^2 x \, dx = \tan x + c$$

To differentiate $y = \sin kx$, where k is a constant, you can define a new variable z by $z = kx$, so that

$$y = \sin kx = \sin z$$

giving $\frac{dy}{dz} = \cos z$

But $\frac{dy}{dx} = \frac{dy}{dz} \times \frac{dz}{dx}$ (function of a function)

and since $z = kx$

then $\frac{dz}{dx} = k$

So $\frac{dy}{dx} = (\cos z) \times k$

giving $\frac{dy}{dx} = k \cos kx$



Exercises

1. Find the derivative of

(a) $\sin 2x$ (b) $\sin \frac{1}{2}x$ (c) $\sin 100x$ (d) $\cos 3x$

2. Find $\frac{d}{dx}(\cos kx)$ and $\frac{d}{dx}(\tan kx)$



Worked Example 2

A spherical balloon is blown up so that its volume increases at a constant rate of $10 \text{ cm}^3/\text{s}$. Find the rate of increase of the radius when the volume of the balloon is 1000 cm^3 .



Solution

Now $v = \frac{4}{3}\pi r^3$

giving $\frac{dv}{dt} = \frac{d}{dr}\left(\frac{4}{3}\pi r^3\right) \frac{dr}{dt}$
 $= 4\pi r^2 \frac{dr}{dt}$

But $\frac{dv}{dt} = 10$

giving $10 = 4\pi r^2 \frac{dr}{dt}$

Thus $\frac{dr}{dt} = \frac{5}{2\pi r^2}$

When the volume is 1000 cm^3 the radius can be calculated from

$$1000 = \frac{4}{3}\pi r^3$$


$$\Rightarrow r = \left(\frac{750}{\pi}\right)^{\frac{1}{3}}$$

$$\Rightarrow \frac{dr}{dt} = \frac{5}{2\pi} \left(\frac{750}{\pi}\right)^{-\frac{2}{3}} \approx 0.021$$



Exercises

- The radius of a circular disc is increasing at a constant rate of 0.003 cm/s . Find the rate at which the area is increasing when the radius is 20 cm .
- A spherical balloon is inflated by gas being pumped at a constant rate of $200 \text{ cm}^3/\text{s}$. What is the rate of increase of the surface area of the balloon when its radius is 100 cm ?
- The volume of water in a container is given by $xe^{-2x} \text{ cm}^3$ where x is the depth of the water in the container. Find the rate of increase of volume when the depth is 0.2 cm and increasing at a rate of 0.1 cm s^{-1} . Also determine when the rate of increase of volume is stationary.
- The area of the region between two concentric circles of radii x and y ($x > y$) is denoted by A . Given that x is increasing at the rate of 3 ms^{-1} , y is increasing at the rate of 4 ms^{-1} , and when $t = 0$, $x = 5 \text{ m}$ and $y = 2 \text{ m}$, find
 - the rate of increase of A when $t = 0$;
 - the ratio of x to y when A begins to decrease;
 - the time at which A is zero.



3 Integration by Substitution

This method is the equivalent to using a change of variable when differentiating composite functions.

In general, if

$$y = \int f(x) dx$$

then $\frac{dy}{dx} = f(x)$

If we let $x = g(u)$ for some function g of a new variable u , then y becomes a function of u and

$$\begin{aligned} \frac{dy}{du} &= \frac{dy}{dx} \times \frac{dx}{du} \\ \Rightarrow \frac{dy}{du} &= f(x) \frac{dx}{du} = f(g(u)) \frac{dx}{du} \\ \Rightarrow y &= \int f(g(u)) \frac{dx}{du} du \end{aligned}$$



Worked Example 1

Find $\int x(3x-1)^{\frac{1}{2}} dx$ using a suitable substitution.



Solution

If $u = 3x - 1$ then $\frac{du}{dx} = 3$ and $x = \frac{u+1}{3}$. Hence the integral becomes

$$\begin{aligned} \int x(3x-1)^{\frac{1}{2}} \frac{dx}{du} du &= \int \left(\frac{u+1}{3}\right) u^{\frac{1}{2}} \frac{du}{3} \\ &= \frac{1}{9} \int \left(u^{\frac{3}{2}} + u^{\frac{1}{2}}\right) du \\ &= \frac{1}{9} \left(\frac{2}{5} u^{\frac{5}{2}} + \frac{2}{3} u^{\frac{3}{2}}\right) + c \\ &= \frac{1}{9} \left(\frac{2}{5} (3x-1)^{\frac{5}{2}} + \frac{2}{3} (3x-1)^{\frac{3}{2}}\right) + c \\ &= \frac{2}{45} (3x-1)^{\frac{5}{2}} + \frac{2}{27} (3x-1)^{\frac{3}{2}} + c \end{aligned}$$



Worked Example 2

Find $\int_0^1 x(3x^2+2)^3 dx$



Solution

You can rewrite this integral using the substitution

$$u = (3x^2 + 2) \Rightarrow \frac{du}{dx} = 6x$$

The limits on u for the integration are determined from

$$x = 0 \Rightarrow u = 2$$

$$x = 1 \Rightarrow u = 5$$

$$\begin{aligned} \text{Hence } \int_0^1 x(3x^2 + 2)^3 dx &= \int_2^5 x(3x^2 + 2)^3 \frac{dx}{du} du \\ &= \int_2^5 x(3x^2 + 2)^3 \frac{1}{6x} du \\ &= \int_2^5 \frac{1}{6} u^3 du \\ &= \left[\frac{1}{24} u^4 \right]_2^5 \\ &= \frac{1}{24} [5^4 - 2^4] \\ &= \frac{1}{24} (625 - 16) \\ &= \frac{609}{24} \end{aligned}$$



Exercises

1. Find the following integrals, using the suggested substitution.

$$(a) \int x(x^2 - 3)^4 dx \quad (u = x^2 - 3)$$

$$(b) \int x\sqrt{1-x^2} dx \quad (u = 1-x^2)$$

$$(c) \int \cos x \sin^4 x dx \quad (u = \sin x)$$

$$(d) \int e^x \sqrt{1+e^x} dx \quad (u = 1+e^x)$$

2. Using the substitution $x = 4\sin^2 \theta$ or otherwise, show that

$$\int_0^2 \sqrt{x(4-x)} dx = \pi$$

3. Using the substitution $x = 3 \tan \theta$, evaluate

$$\int_0^3 \frac{1}{(9+x^2)^2} dx$$

4 Integration by Parts

Our second method of integration is a very important one which is particularly useful for a number of different integrals. You can see how it works by first differentiating with respect to x ,

$$x \sin x + \cos x$$

Now use your answer to find the integral of $x \cos x$.

What form will the integral of $x \sin x$ take?

Check your answers by differentiating.



Worked Example 1

By assuming

$$\int x e^x dx = (ax + b)e^x + c$$

find the values of the constants a and b so that the result is valid.



Solution

We can find a and b by differentiating both sides, that is

$$\begin{aligned} x e^x &= \frac{d}{dx} ((ax + b)e^x + c) \\ &= (ax + b)e^x + ae^x + 0 \\ &= (ax + a + b)e^x \end{aligned}$$

So $a = 1$ and $a + b = 0$

giving $b = -1$

Hence

$$\int x e^x dx = (x - 1)e^x + c$$

By now you should be wondering if there is a more precise method of evaluating integrals of the type above, where the method requires inspired guesswork to start with. In fact, there is a more formal method.

If u and v are two functions of x then from your work on differentiation you know that

$$\frac{d}{dx}(uv) = v \frac{du}{dx} + u \frac{dv}{dx}$$

and integrating each side with respect to x gives

$$uv = \int v \frac{du}{dx} dx + \int u \frac{dv}{dx} dx$$

$$\Rightarrow \int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$$



Worked Example 2

Find $\int x \cos x \, dx$



Solution

Here $u = x$ and $\frac{dv}{dx} = \cos x$, so $v = \sin x$. This gives

$$\begin{aligned}\int x \cos x \, dx &= x \sin x - \int \sin x \cdot 1 \, dx \\ &= x \sin x + \cos x + c\end{aligned}$$

We can use the same method to find the integral in Worked Example 1.



Worked Example 3

Find $\int x e^x \, dx$



Solution

Here $u = x$, $\frac{dv}{dx} = e^x$, giving $v = e^x$ and

$$\begin{aligned}\int x e^x \, dx &= x e^x - \int e^x \cdot 1 \, dx \\ &= x e^x - e^x + c \\ &= e^x(x-1) + c\end{aligned}$$

One of the difficulties with integration by parts is that of identifying u and $\frac{dv}{dx}$. For example, if

$$I = \int x \ln x \, dx$$

then taking $u = x, \frac{dv}{dx} = \ln x$ will not really work as you might not be able to solve for v .

But taking

$$u = \ln x, \frac{dv}{dx} = x$$

gives $v = \frac{1}{2}x^2$, and

$$\begin{aligned}I &= \frac{1}{2}x^2 \ln x - \int \left(\frac{1}{2}x^2\right) \frac{1}{x} \, dx \\ &= \frac{1}{2}x^2 \ln x - \int \frac{1}{2}x \, dx \\ &= \frac{1}{2}x^2 \ln x - \frac{1}{4}x^2 + c\end{aligned}$$

It is also important to note that some functions which cannot be immediately integrated can be dealt with by writing the integrand as $1 \times (\text{function})$ since 1 can be integrated. This is illustrated below.



Worked Example 4

Find $\int \ln x \, dx$



Solution

$$I = \int \ln x \, dx = \int 1 \times \ln x \, dx$$

So take $u = \ln x$, $\frac{dv}{dx} = 1$ to give $v = x$ and

$$\begin{aligned} I &= x \ln x - \int x \cdot \frac{1}{x} \, dx \\ &= x \ln x - \int 1 \, dx \\ &= x \ln x - x + c \end{aligned}$$

Finally in this section it should be noted that we can have limits on integration in the usual way, for example

$$\begin{aligned} \int_0^{\frac{\pi}{2}} x \sin x \, dx &= [-x \cos x]_0^{\frac{\pi}{2}} - \int_0^{\frac{\pi}{2}} 1(-\cos x) \, dx \\ &= (0 - 0) + \int_0^{\frac{\pi}{2}} \cos x \, dx \\ &= [\sin x]_0^{\frac{\pi}{2}} \\ &= 1 \end{aligned}$$



Exercises

1. Find $\int x^2 \ln x \, dx$.
2. By writing $\cos^3 \theta$ as $\cos^2 \theta \cos \theta$ find

$$\int \cos^3 \theta \, d\theta$$

and hence show that

$$\int_0^x \cos^3 \theta \, d\theta = 0.$$

3. Evaluate $\int_0^{\frac{\pi}{2}} \sin^3 x \, dx$.
4. Find $\int x^3 \ln(4x) \, dx$.

5. Use integration by parts twice to find

$$\int x^2 \cos 2x \, dx$$

6. Given that $\int x \cos x \, dx = x \sin x + \cos x + c$ show that

$$\int x^2 \sin x \, dx = x^2 \cos x + 2x \sin x + 2 \cos x + k$$

7. Given that $\int x e^x \, dx = x e^x - e^x + c$ find

(a) $\int x^2 e^x \, dx$ (b) $\int x^2 e^{-x} \, dx$

5 First Order Differential Equations

You might well wonder why you have studied integration in such depth. The main reason is the crucial role it plays in solving what are known as **differential equations**. Here you will only be dealing with first order differential equations, which can be expressed, in general, in the form

$$\frac{dy}{dx} = f(x, y)$$

when $y = y(x)$ is an unknown function of x , and f is a given function of x and y .

Examples of such equations include

$$\frac{dy}{dx} = xy$$

$$\frac{dy}{dx} = x + y$$

$$\frac{dy}{dx} = \frac{x}{y} \quad (y \neq 0)$$

but first you will see how they are formed.

As an example of the process, consider the problem of modelling the way in which human populations change.

An English economist, *Thomas Malthus*, in his 1802 article "An Essay on the Principle of Population" formulated the first model which attempted to describe and predict the way in which the magnitude of the human population was changing. In mathematical terms you can summarise his assumptions by introducing the variable $N = N(t)$ to represent the total population, where t is time. In a small time interval, say δt , Malthus argued that both births and deaths are proportional to the population size and the time interval. Hence in time δt , there will be $(\alpha N \delta t)$ births and $(\beta N \delta t)$ deaths where α, β are positive constants, and so the increase in the population size is given by

$$\delta N = \alpha N \delta t - \beta N \delta t = (\alpha - \beta) N \delta t$$

Writing $\alpha - \beta = \gamma$, and dividing by δt gives

$$\frac{\delta N}{\delta t} = \gamma N$$

Taking the limit as $\delta t \rightarrow 0$ results in the differential equation

$$\boxed{\frac{dN}{dt} = \gamma N}$$

Added to this is an initial condition; for example, $N = N_0$ at $t = 0$.

This is an example of a 'variable separable' differential equation, since it can be written in the form

$$\frac{1}{N} \frac{dN}{dt} = \gamma$$

and integration with respect to t ,

$$\int \frac{1}{N} \frac{dN}{dt} dt = \int \gamma dt$$

$$\int \frac{1}{N} dN = \int \gamma dt$$

$$\Rightarrow \ln N = \gamma t + c \quad (c \text{ constant})$$

$$\begin{aligned} \Rightarrow N &= e^{\gamma t + c} \\ &= e^{\gamma t} e^c \end{aligned}$$

$$\Rightarrow N = Ke^{\gamma t} \quad (K = e^c)$$

But

$$N = N_0 \text{ at } t = 0$$


giving

$$N_0 = Ke^0 = K$$

Hence the complete solution is

$$\boxed{N = N_0 e^{\gamma t}}$$

This solution is illustrated opposite for positive γ .



You can see how to use this population model by applying it to the 1790 and 1800 USA population figures shown opposite.

The time $t = 0$ corresponds to 1790, and $t = 1$ to 1800 etc; so

$$N_0 = 3.9$$

Year	USA Population (in millions)
1790	3.9
1800	5.3

The model solution is

$$N(t) = 3.9e^{\gamma t}$$

and γ is determined by the population value at $t = 1$ (1800). This gives

$$5.3 = 3.9e^{\gamma}$$

$$\Rightarrow e^{\gamma} = \frac{5.3}{3.9}$$

$$\Rightarrow \gamma = \ln\left(\frac{5.3}{3.9}\right) \approx 0.3067$$

and $N(t) = 3.9e^{(0.3067)t}$



Worked Example 1

Use the model above to predict the USA population for 1810 ($t = 2$) to 1930 ($t = 14$). Compare your prediction with the real values given opposite.



Solution

$$\begin{aligned} t = 2 \Rightarrow N(2) &= 3.9e^{0.3067 \times 2} \\ &= 3.9e^{0.6134} \\ &\approx 7.2 \end{aligned}$$

$$\begin{aligned} t = 14 \Rightarrow N(14) &= 3.9e^{0.3067 \times 14} \\ &\approx 285.6 \end{aligned}$$

So the model give an accurate prediction for 1810 but is much too high for 1930.

The model gives a reasonably good fit for quite some time, but eventually predicts a population growing far too quickly.

Why does the model eventually go wrong?

Let us go back to the original assumption and consider what factors have been neglected. The model as it stands predicts the growth of a population in an ideal situation where there are no limits (e.g. food, land, energy) to continued growth. In 1837, Verhulst proposed an extension to the Malthusian model which took into account the 'crowding' factor. In his model, he assumed that the population change was proportional to

- (i) the current population level N
- (ii) the ratio of unused population resources to total population resources when it is assumed that a maximum population size of N_{∞} can be sustained,

i.e.
$$\frac{(N_{\infty} - N)}{N_{\infty}}$$

Year	USA Population (in millions)
1810	7.2
1820	9.6
1830	12.9
1840	17.1
1850	23.2
1860	31.4
1870	38.6
1880	50.2
1890	62.9
1900	76.0
1910	92.0
1920	106.5
1930	123.2

Thus

$$\frac{dN}{dt} = \gamma N \frac{(N_\infty - N)}{N_\infty}$$

i.e.

$$\boxed{\frac{dN}{dt} = \gamma N - \frac{\gamma N^2}{N_\infty}}$$

When N is small, γN will be the dominant term on the right hand side, and so we are back with the Malthusian model; but as N becomes large, the $\frac{-\gamma N^2}{N_\infty}$ term becomes important.

This is again a first order differential equation, but much more difficult to solve.

It can in fact be solved and has solution

$$N(t) = \frac{N_\infty}{\left\{1 + \left[\frac{N_\infty}{N_0} - 1\right] e^{-\gamma t}\right\}}$$

where $N = N_0$ at $t = 0$.

As $t \rightarrow \infty$, you can see that $N \rightarrow N_\infty$

Returning to simple first order differential equations, you will see how to solve them in the following two examples.



Worked Example 2

For $x \geq 0$, solve $\frac{dy}{dx} = \frac{x}{y}$ given that $y = 1$ when $x = 0$.



Solution

You can write this as

$$y \frac{dy}{dx} = x$$

and integrating both sides with respect to x .

$$\begin{aligned} \int y \frac{dy}{dx} dx &= \int x dx \\ \Rightarrow \int y dy &= \int x dx \\ \Rightarrow \frac{1}{2} y^2 &= \frac{1}{2} x^2 + K \quad (K \text{ constant}) \end{aligned}$$

Since $y = 1$ when $x = 0$,

$$\frac{1}{2} = 0 + K$$

$$\Rightarrow K = \frac{1}{2}$$

and

$$y^2 = x^2 + 1$$

$$\Rightarrow y = \sqrt{1 + x^2}$$



Worked Example 3

Solve $\frac{dy}{dx} = \frac{y}{x}$ given that $y = 1$ when $x = 1$.



Solution

As before

$$\frac{1}{y} \frac{dy}{dx} = \frac{1}{x}$$

$$\Rightarrow \int \frac{1}{y} \frac{dy}{dx} dx = \int \frac{1}{x} dx$$

$$\Rightarrow \int \frac{1}{y} dy = \int \frac{1}{x} dx$$

$$\Rightarrow \ln y = \ln x + C \quad (C \text{ constant})$$

Writing $C = \ln K$,

$$\ln y = \ln x + \ln K$$

$$\Rightarrow \ln y = \ln(Kx)$$

$$\Rightarrow y = Kx$$

Since $y = 1$ when $x = 1$,

$$1 = K \cdot 1$$

giving

$$K = 1$$

and the solution

$$y = x$$



Exercises

Solve the following first order differential equations.

1. $3y \frac{dy}{dx} = 5x^2$ given that $y = 1$ when $x = 0$

2. $x \frac{dy}{dx} = 3y$ given that $y = 2$ when $x = 1$

3. $\frac{dy}{dx} = y^2$ given that $y = 1$ when $x = 1$

4. $\frac{1}{x} \frac{dy}{dx} = \frac{1}{(x^2 + 1)}$ given that $y = 0$ when $x = 0$

5. $e^x \frac{dy}{dx} = \frac{x}{y}$

6. $\frac{dy}{dx} = \frac{y+1}{x+2}$

7. $x \frac{dy}{dx} = \frac{1}{y} + y$

8. During the initial stages of the growth of yeast cells in a culture, the rate of increase in the number of cells is proportional to the number of cells present.

If $n = n(t)$ represents the number of cells at time t , show that

$$\frac{dn}{dt} = kn$$

for some constant k . If the cells double in unit time, determine k , and solve for n in terms of t and the initial number of cells, n_0 .